

ACCESSION NUMBER: 2000:455053 HCAPLUS
DOCUMENT NUMBER: 134:114464
TITLE: Synthetic peptides that inhibit binding of the collagen type II 261-273 epitope to rheumatoid arthritis-associated HLA-DR1 and -DR4 molecules and collagen-specific T-cell responses
AUTHOR(S): Fridkis-Hareli, M.; Rosloniec, E. F.; Fugger, L.; Strominger, J. L.
CORPORATE SOURCE: Department of Clinical Immunology, Aarhus University Hospital, Aarhus, Den.
SOURCE: Hum. Immunol. (2000), 61(7), 640-650
CODEN: HUIMDQ; ISSN: 0198-8859
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Copolymer 1 [Cop 1, poly (Y, E, A, K)] is a random synthetic amino acid copolymer effective in the treatment of relapsing forms of multiple sclerosis (MS), a disease that is linked to HLA-DR2 (DRB1*1501). Another copolymer [poly (Y, A, K)] was also identified that binds to rheumatoid arthritis (RA)-assocd. HLA-DR1 (DRB1*0101) or HLA-DR4 (DRB1*0401) mols. and inhibits the response of HLA-DR1- and -DR4-restricted T cell clones to an immunodominant epitope of collagen type II (CII) 261-273 (a candidate autoantigen in RA). In the present study various peptides have been synthesized based on binding "motifs" of Cop 1 for HLA-DR1 and -DR4 mols. Those peptides with K at P-1 or K at P8 were particularly effective as inhibitors of binding of CII 261-273, of Cop 1 and of the influenza virus hemagglutinin peptide 306-318 to these class II proteins. Moreover, several of them were also potent inhibitors of the CII 261-273-reactive T cell clones. These findings suggest that small peptides or their more stable derivs. may be able to substitute for copolymers in the treatment of RA, and by implication of MS.

REFERENCE COUNT: 63

REFERENCE(S): (1) Aharoni, R; Eur J Immunol 1993, V23, P17 HCAPLUS
(2) Andersson, E; Proc Natl Acad Sci USA 1998, V95, P7574 HCAPLUS
(4) Brand, D; J Immunol 1994, V152, P3088 HCAPLUS
(5) Brown, J; Nature 1993, V364, P33 HCAPLUS
(6) Busch, R; J Immunol 1991, V147, P1292 HCAPLUS

ACCESSION NUMBER: 2000:455053 HCAPLUS
DOCUMENT NUMBER: 134:114464
TITLE: Synthetic peptides that inhibit binding of the collagen type II 261-273 epitope to rheumatoid arthritis-associated HLA-DR1 and -DR4 molecules and collagen-specific T-cell responses
AUTHOR(S): Fridkis-Hareli, M.; Rosloniec, E. F.; Fugger, L.; Strominger, J. L.
CORPORATE SOURCE: Department of Clinical Immunology, Aarhus University Hospital, Aarhus, Den.
SOURCE: Hum. Immunol. (2000), 61(7), 640-650
CODEN: HUIMDQ; ISSN: 0198-8859
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Copolymer 1 [Cop 1, poly (Y, E, A, K)] is a random synthetic amino acid copolymer effective in the treatment of relapsing forms of multiple sclerosis (MS), a disease that is linked to HLA-DR2 (DRB1*1501). Another copolymer [poly (Y, A, K)] was also identified that binds to rheumatoid arthritis (RA)-assocd. HLA-DR1 (DRB1*0101) or HLA-DR4 (DRB1*0401) mols. and inhibits the response of HLA-DR1- and -DR4-restricted T cell clones to an immunodominant epitope of collagen type II (CII) 261-273 (a candidate autoantigen in RA). In the present study various peptides have been synthesized based on binding "motifs" of Cop 1 for HLA-DR1 and -DR4 mols. Those peptides with K at P-1 or K at P8 were particularly effective as inhibitors of binding of CII 261-273, of Cop 1 and of the influenza virus hemagglutinin peptide 306-318 to these class II proteins. Moreover, several of them were also potent inhibitors of the CII 261-273-reactive T cell clones. These findings suggest that small peptides or their more stable derivs. may be able to substitute for copolymers in the treatment of RA, and by implication of MS.

REFERENCE COUNT: 63

REFERENCE(S): (1) Aharoni, R; Eur J Immunol 1993, V23, P17 HCAPLUS
(2) Andersson, E; Proc Natl Acad Sci USA 1998, V95, P7574 HCAPLUS
(4) Brand, D; J Immunol 1994, V152, P3088 HCAPLUS
(5) Brown, J; Nature 1993, V364, P33 HCAPLUS
(6) Busch, R; J Immunol 1991, V147, P1292 HCAPLUS

ACCESSION NUMBER: 2000:455053 HCAPLUS
DOCUMENT NUMBER: 134:114464
TITLE: Synthetic peptides that inhibit binding of the collagen type II 261-273 epitope to rheumatoid arthritis-associated HLA-DR1 and -DR4 molecules and collagen-specific T-cell responses
AUTHOR(S): Fridkis-Hareli, M.; Rosloniec, E. F.; Fugger, L.; Strominger, J. L.
CORPORATE SOURCE: Department of Clinical Immunology, Aarhus University Hospital, Aarhus, Den.
SOURCE: Hum. Immunol. (2000), 61(7), 640-650
CODEN: HUIMDQ; ISSN: 0198-8859
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Copolymer 1 [Cop 1, poly (Y, E, A, K)] is a random synthetic amino acid copolymer effective in the treatment of relapsing forms of multiple sclerosis (MS), a disease that is linked to HLA-DR2 (DRB1*1501). Another copolymer [poly (Y, A, K)] was also identified that binds to rheumatoid arthritis (RA)-assocd. HLA-DR1 (DRB1*0101) or HLA-DR4 (DRB1*0401) mols. and inhibits the response of HLA-DR1- and -DR4-restricted T cell clones to an immunodominant epitope of collagen type II (CII) 261-273 (a candidate autoantigen in RA). In the present study various peptides have been synthesized based on binding "motifs" of Cop 1 for HLA-DR1 and -DR4 mols. Those peptides with K at P-1 or K at P8 were particularly effective as inhibitors of binding of CII 261-273, of Cop 1 and of the influenza virus hemagglutinin peptide 306-318 to these class II proteins. Moreover, several of them were also potent inhibitors of the CII 261-273-reactive T cell clones. These findings suggest that small peptides or their more stable derivs. may be able to substitute for copolymers in the treatment of RA, and by implication of MS.

REFERENCE COUNT: 63

REFERENCE(S): (1) Aharoni, R; Eur J Immunol 1993, V23, P17 HCAPLUS
(2) Andersson, E; Proc Natl Acad Sci USA 1998, V95, P7574 HCAPLUS
(4) Brand, D; J Immunol 1994, V152, P3088 HCAPLUS
(5) Brown, J; Nature 1993, V364, P33 HCAPLUS
(6) Busch, R; J Immunol 1991, V147, P1292 HCAPLUS